



Outline

- Sediment Investigation
 - Geophysical Investigation
 - High Resolution Core Site Selection
 - Core Processing
- Water Column Investigation
 - SPMDs
 - Large volume filtration
 - High flow event
- What's Next



Sediment Investigation


- Geophysical Results
 - Sediment Texture Analysis
 - Geophysical Mapping
 - Extent of Sediment Types




Sediment Texture Analyses


- Grain size distribution vs field classification
- Grain size distribution vs Side Scan
 Sonar interpretation
- Field classification vs SSS interpretation













Side Scan Sonar Observations

- Preliminary sediment assignments agree well
 - About 70 percent accuracy
 - Probably 3 or 4 true classes
- SSS maps provide a basis to map fine sediments

Maps of observations



Comparison of SSS Results and Deposition Rates



Average Rate of Depth Change (inches / year)

Deposition

Rate

Legend

< -5

-5 - -4

-4 - -3

-3 - -2

-2 - -1

-1 - 0

0 - 1

1-2

2 - 3

3 - 4

4 - 5

5 +

Sediment **Texture** Legend

■ Silt

□ Silt and Sand

Sand


■ Sand and Gravel

■ Rock and Coarse Gravel


Maps


- RM 2-3 Deposition
- RM 2-3 Sediment Texture
- RM 12-13 Deposition
- RM 12-13 Sediment Texture





Sediment Texture by River Mile






- Silt
- ☐ Silt and Sand
- Sand
- Sand and Gravel
- Rock and Coarse Grave




Deposition Rate vs Texture, by Area

Sonar Classification



Correlation of Sediment Type and Average Annual Deposition Rate



High Resolution Core Site Selection

Sediment Deposition Rate Maps
Sediment Texture Maps
Historical Data



Site Selection




Be-7 Survey


- Survey conducted in early September immediately following receipt of sediment texture maps.
- Achieved 1 week turn-around on samples to enable final core site selection.
- Values as high as 9 pCi/g



Surface Be⁷ vs River Mile



Surface Cs¹³⁷ vs River Mile








High Resolution Core Site Selection

- Be7-bearing fine sediment identified in many locations
- Core sites available every 2 miles or so to RM 12.
 - Nothing available between RM 12 and 17
- 15 core sites planned from baseline area to Newark Bay entrance.








High Resolution Core Collection

- Obtained 3 to 4 cores per day
- 4 to 5 cores per location
- 1 to 2 cores for sectioning
- Processed 44 to 88 sections per day
- Processed, labeled, sealed, recorded, stored and/or shipped up to 300+ sample jars per day
- 14 cores successfully obtained
- Sample management via radionuclide screening and sample preservation to reduce costs

















INDEPENDENT ENVIRONMENTAL ENGINEERS, SCIENTISTS AND CONSULTANTS



INDEPENDENT ENVIRONMENTAL ENGINEERS, SCIENTISTS AND CONSULTANTS

Semi-Permeable Membrane Devices (SPMD)

- Total of 2 deployments, 8 stations each
 - 4 tributaries
 - 4 main stem stations
- 1st deployment Week of Sept 19th, 2005
- Second deployment in this week.
- Each SPMD canister contains 5 lipid bags to provide enough volume for analysis of : PCB Congeners, Dioxin/furans, PAHs, Pesticides.














Large Volume Filtration Samples

- Two events, 3 stations per event
 - September
 - November
- Six samples/station
 - Large volume dissolved and suspended matter via Infiltrex
 - Large volume dissolved and suspended matter via TOPS (dissolved sample shipped)
 - Large volume whole water (lab to filter and analyze separate phases)
- September event demonstrated need to filter to 0.5 um



















What's Next

- High Resolution Cores
 - Complete radionuclide analysis/interpretation
 - Identify cores and samples for further analysis.
 - First samples should ship by Dec 1
 - Data available in spring 2006
- SPMDs
 - Samples retrieved this week
 - Second deployment begins this week
 - Data available this winter
- Large Volume Samples
 - Second round later this month
 - Samples from first event undergoing analysis
- Small Volume Composites
 - Sampling event next Tuesday (flood tide)
 - Ebb tide event to be scheduled in next 2 to 4 weeks

